

Phenomenological approach to symmetry breaking pattern of democratic mass matrix

Junpei Harada*

Department of Physics, Hiroshima University, Hiroshima 739-8526, Japan

Abstract

We investigate the symmetry breaking pattern of the democratic mass matrix model, which leads to the small flavor mixing in quark sector and bi-large mixing in lepton sector. We present the symmetry breaking matrices in quark sector which are determined by alternative ways instead of conventional ansatz. These matrices might be useful for understanding the origin of democratic symmetry and its breaking.

*e-mail: harada@theo.phys.sci.hiroshima-u.ac.jp

1 Introduction

One of the most important problems in particle physics is to understand the flavor structure of quarks and leptons. Since the feature of the flavor structure appear in the quark-lepton mass matrix, it is important to investigate the quark-lepton mass matrix for understanding the flavor structure.

Various phenomenological mass matrix models of quarks and leptons are proposed. One of the most attractive mass matrix models is democratic mass matrix model [1], which has $S_{3L} \times S_{3R}$ permutation symmetry (democratic symmetry). This symmetry means that right-handed fermion and left-handed fermion independently have the permutation symmetry among three generations. This democratic mass matrix model induces the small mixing angles in quark sector and bi-large mixing angles in lepton sector [2][3]. The democratic mass matrix model also induces small mixing angle U_{e3} in lepton sector, which is consistent with the experimental bound from CHOOZ reactor experiment [4]. Thus the democratic mass matrix model is very successful phenomenologically.

In spite of these successes, however, the democratic mass matrix induces the massless quarks and charged leptons of the first and the second generations. Then Cabibbo-Kobayashi-Maskawa (CKM) quark flavor mixing matrix is exactly equals to a unit matrix in the democratic symmetry limit. To acquire realistic fermion masses and mixing angles, democratic symmetry must be broken and small symmetry breaking terms are required. If we know the mechanism of democratic symmetry breaking, we can predict the quark-lepton mass spectrum and all mixing angles. Unfortunately, however, such symmetry breaking mechanism is not known. In the literature, some works have assumed these small symmetry breaking terms and given some predictions. This approach has been known very successful for quark sector [5][6] and lepton sector [7][8][9]. However, there are possibilities that other ansatz can also account for the observed masses and mixing angles. Then it is very important to understand the mechanism of democratic symmetry breaking.

In this letter, we present the symmetry breaking terms which are determined by alternative ways instead of conventional ansatz. These symmetry breaking terms might be useful for understanding the origin of democratic symmetry and the mechanism of its breaking. Although we are especially interested in the lepton sector, we investigate the symmetry breaking terms of quark sector in this letter. The reasons are as follows. First, the experimental information of quark sector is much rather than that of lepton

sector. Second, since both up-type quarks and down-type quarks are Dirac particles, they are treated in an equal footing in the democratic mass matrix model. Third, we expect that the symmetry breaking pattern of down quarks can be applied for charged leptons.

This letter is organized as follows. In sec. 2 we briefly review the democratic mass matrix model. In sec. 3 we present the symmetry breaking terms of quark sector. In sec. 4 we comment on lepton sector. Conclusion is denoted in sec. 5.

2 Democratic Mass Matrix

We briefly review the democratic mass matrix model. At first, we concentrate on quark sector. The democratic mass matrix has $S_{3L} \times S_{3R}$ permutation symmetry, i.e.,

$$M_u^{[0]} = \frac{m_t}{3} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}, \quad (2.1)$$

where m_t is top quark mass. The down quark mass matrix has the same structure. This matrix is a unique representation of the $S_{3L} \times S_{3R}$ symmetric matrix. This matrix is diagonalized as

$$V_u^{[0]\dagger} M_u^{[0]} V_u^{[0]} = M_u^{[0]diag} = diag(0, 0, m_t). \quad (2.2)$$

The unitary matrix $V_u^{[0]}$ that diagonalizes the democratic mass matrix $M_u^{[0]}$ is

$$V_u^{[0]} = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{6} & 1/\sqrt{3} \\ -1/\sqrt{2} & 1/\sqrt{6} & 1/\sqrt{3} \\ 0 & -2/\sqrt{6} & 1/\sqrt{3} \end{bmatrix}. \quad (2.3)$$

Since there is no difference between up quark mass matrix and down quark mass matrix(except for $m_t \neq m_b$), unitary matrices that diagonalizes mass matrix of up quarks and one of down quarks are same each other, i.e., $V_u^{[0]} = V_d^{[0]}$. Then Cabbibo-Kobayashi-Maskawa(CKM) quark flavor mixing matrix is predicted as

$$V_{CKM}^{[0]} = V_u^{[0]\dagger} V_d^{[0]} = diag(1, 1, 1). \quad (2.4)$$

From experimental observations, it is well known that

$$M_q^{diag} \simeq diag(\epsilon_q^4, \epsilon_q^2, 1)m_{t/b}, \quad (2.5)$$

where $q = u$ or d , and CKM mixing matrix in Wolfenstein parametrization is given by

$$V_{CKM} = \begin{bmatrix} 1 - \frac{\lambda^2}{2} & \lambda & 0 \\ -\lambda & 1 - \frac{\lambda^2}{2} & A\lambda^2 \\ 0 & -A\lambda^2 & 1 \end{bmatrix} + O(\lambda^3). \quad (2.6)$$

Thus, in quark sector, the predictions of the democratic mass matrix model are completely consistent with experimental data with appropriate approximations, i.e.,

$$M_u^{diag} = diag(0, 0, m_t) + O(\epsilon_u^2), \quad (2.7)$$

$$M_d^{diag} = diag(0, 0, m_b) + O(\epsilon_d^2), \quad (2.8)$$

$$V_{CKM} = diag(1, 1, 1) + O(\lambda). \quad (2.9)$$

Let's turn to the lepton sector. Since the charged leptons are Dirac particles, the charged lepton mass matrix has the same structure of quark mass matrix. Assuming that the neutrinos are the Majorana type, there are two independent mass matrices that are invariant under S_{3L} permutation symmetry [7]. Then democratic neutrino mass matrix is given as follows;

$$M_\nu^{[0]} = m_\nu \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} + m_\nu r \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}, \quad (2.10)$$

where r is arbitrary parameter. Here we take $r = 0$, since it automatically leads to the bi-large mixing angles of solar and atmospheric neutrinos *and* small U_{e3} (We comment on this choice in sec. 4). Thus in the democratic mass matrix model, neutrino masses of three generations are degenerate. Then Maki-Nakagawa-Sakata(MNS) lepton flavor mixing matrix is given by

$$U_{MNS}^{[0]} = V_l^{[0]\dagger} = \begin{bmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 1/\sqrt{6} & 1/\sqrt{6} & -2/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} \end{bmatrix}, \quad (2.11)$$

where $V_l^{[0]}$ is the unitary matrix that diagonalizes the charge lepton mass matrix. The predictions of mixing angles are

$$\sin^2 2\theta_{sol} = 1, \quad \sin^2 2\theta_{atm} = \frac{8}{9}, \quad U_{e3} = 0. \quad (2.12)$$

Thus, in lepton sector, the predictions of democratic mass matrix model are consistent with experimental data with appropriate approximations.

In summary, the democratic mass matrix model is phenomenologically very successful in both quark sector and lepton sector.

3 Symmetry Breaking Matrix

In previous section, we saw the democratic mass matrix is good candidate phenomenologically. However, the democratic mass matrix predict the massless quarks and charged leptons of the first and the second generations. It also predicts that CKM mixing matrix equals to a unit matrix. To aquire realistic fermion masses and mixing angles, small symmetry breaking terms are required. In this section, we present such symmetry breaking terms. We concentrate on the quark sector as denoted in the introduction.

We determine the symmetry breaking terms order by order. For example, V_{ub} equals zero in this letter. If one want to induce realistic V_{ub} , higher order correction terms are required. This point is crucial in our approach.

At first, we consider the symmetry breaking terms which leads to realistic quark masses, while CKM mixing matrix remains a unit matrix (We consider the effects of deviation from a unit matrix later). In this letter, we assume that up quark mass matrix and down quark mass matrix are symmetric mass matrices because of democratic principle. We denote the diagonal mass matrix M_q^{diag} as follows,

$$M_q^{diag} \equiv M_q^{[0]diag} + M_q^{[1]diag} + M_q^{[2]diag}, \quad (3.1)$$

$$= diag(0, 0, 1)m_{t/b} + diag(0, \epsilon_q^2, 0)m_{t/b} + diag(\epsilon_q^4, 0, 0)m_{t/b}. \quad (3.2)$$

Since the unitary matrix $V_q^{[0]}$ that diagonalizes the democratic mass matrix $M_q^{[0]}$ is already known, as long as CKM mixing matrix remains a unit matrix, we can *uniquely* determine the symmetry breaking terms which lead to the massive quarks of the first and the second generations,

$$\begin{aligned} M_q &= V_q^{[0]} M_q^{diag} V_q^{[0]\dagger}, \\ &= \sum_{i=0}^2 V_q^{[i]} M_q^{[i]diag} V_q^{[i]\dagger}, \\ &\equiv M_q^{[0]} + M_q^{[1]} + M_q^{[2]}, \\ &= \frac{m_{t/b}}{3} \left(\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} + \epsilon_q^2 \begin{bmatrix} 1/2 & 1/2 & -1 \\ 1/2 & 1/2 & -1 \\ -1 & -1 & 2 \end{bmatrix} + \epsilon_q^4 \begin{bmatrix} 3/2 & -3/2 & 0 \\ -3/2 & 3/2 & 0 \\ 0 & 0 & 0 \end{bmatrix} \right) \end{aligned} \quad (3.3)$$

Note here that the first order symmetry breaking term, which leads to the charm/strange quark masses, automatically has $S_{2L} \times S_{2R}$ permutaion symmetry [10].

Next we consider the symmetry breaking terms which lead to realistic quark flavor mixing angles. Here we assume that M_u and M_d are the fuction of ϵ_u and ϵ_d , respectively.

We take CKM mixing matrix in $O(\lambda^2)$ approximation, i.e.,

$$V_{CKM} = \begin{bmatrix} 1 & \lambda & 0 \\ -\lambda & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} + O(\lambda^2). \quad (3.4)$$

Note here that CP is conserved at this order since CP phase exits in $O(\lambda^3)$ terms in Wolfenstein parametrization. We denote the unitary matrix that diagonalizes up/down mass matrix as follows,

$$V_q = V_q^{[0]} + V_q^{[1]} + \dots, \quad (3.5)$$

where $V_q^{[1]}, \dots$ are small correction terms which depend on ϵ_q . Then CKM mixing matrix is given by

$$V_{CKM} = V_u^{[0]\dagger} V_d^{[0]} + V_u^{[1]\dagger} V_d^{[0]} + V_u^{[0]\dagger} V_d^{[1]} + V_u^{[1]\dagger} V_d^{[1]} + \dots. \quad (3.6)$$

The first term obviously equals to a unit matrix, which is zero-th order CKM mixing matrix $V_{CKM}^{[0]}$. Since the fourth term is higher order rather than the second and the third terms, we assume that the second and third terms represent $O(\lambda)$ terms in CKM matrix,

$$V_{CKM}^{[1]} \equiv \begin{bmatrix} 0 & \lambda & 0 \\ -\lambda & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = V_u^{[1]\dagger} V_d^{[0]} + V_u^{[0]\dagger} V_d^{[1]}. \quad (3.7)$$

This equation should hold for arbitrary ϵ_q and we can derive the first order correction term $V_q^{[1]}$,

$$V_q^{[1]} = a_q \begin{bmatrix} -1/\sqrt{6} & 1/\sqrt{2} & 0 \\ -1/\sqrt{6} & -1/\sqrt{2} & 0 \\ 2/\sqrt{6} & 0 & 0 \end{bmatrix}, \quad (3.8)$$

where $\sin \theta_{12}^q \equiv \lambda = a_d - a_u$. Note here that parameter a_q depends on ϵ_q . Using this symmetry breaking terms of unitary matrix, we can determine the symmetry breaking mass matrix which induces $O(\lambda)$ terms in CKM mixing matrix,

$$\begin{aligned} M_q &= V_q M_q^{diag} V_q^\dagger, \\ &= \sum_{i=0}^2 V_q M_q^{[i]diag} V_q^\dagger, \\ &= \sum_{i=0}^2 \left(V_q^{[0]} M_q^{[i]diag} V_q^{[0]\dagger} + V_q^{[1]} M_q^{[i]diag} V_q^{[0]\dagger} + V_q^{[0]} M_q^{[i]diag} V_q^{[1]\dagger} + \dots \right). \end{aligned} \quad (3.9)$$

The first term represents eq. (3.3). The second and third terms which induces $O(\lambda)$ terms in CKM mixing matrix are

$$\sum_{i=0}^2 \left(V_q^{[1]} M_q^{[i]diag} V_q^{[0]\dagger} + V_q^{[0]} M_q^{[i]diag} V_q^{[1]\dagger} \right) = \frac{m_{t/b}}{3} \sqrt{3} a_q (\epsilon_q^2 - \epsilon_q^4) \begin{bmatrix} 1 & 0 & -1 \\ 0 & -1 & 1 \\ -1 & 1 & 0 \end{bmatrix}. \quad (3.10)$$

In summary, the democratic quark mass matrix with small symmetry breaking terms which lead to the realistic quark masses and CKM mixing matrix in $O(\lambda^2)$ approximation is given by

$$\begin{aligned} M_q = & \frac{m_{t/b}}{3} \left(\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} + \epsilon_q^2 \begin{bmatrix} 1/2 & 1/2 & -1 \\ 1/2 & 1/2 & -1 \\ -1 & -1 & 2 \end{bmatrix} + \epsilon_q^4 \begin{bmatrix} 3/2 & -3/2 & 0 \\ -3/2 & 3/2 & 0 \\ 0 & 0 & 0 \end{bmatrix} \right) \\ & + \frac{m_{t/b}}{3} \left(\sqrt{3} a_q (\epsilon_q^2 - \epsilon_q^4) \begin{bmatrix} 1 & 0 & -1 \\ 0 & -1 & 1 \\ -1 & 1 & 0 \end{bmatrix} \right). \end{aligned} \quad (3.11)$$

This mass matrix is diagonalized as

$$M_q^{diag} = V_q^\dagger M_q V_q = diag(\epsilon_q^4, \epsilon_q^2, 1) m_{t/b}, \quad (3.12)$$

where unitary matrix V_q is

$$V_q = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{6} & 1/\sqrt{3} \\ -1/\sqrt{2} & 1/\sqrt{6} & 1/\sqrt{3} \\ 0 & -2/\sqrt{6} & 1/\sqrt{3} \end{bmatrix} + a_q \begin{bmatrix} -1/\sqrt{6} & 1/\sqrt{2} & 0 \\ -1/\sqrt{6} & -1/\sqrt{2} & 0 \\ 2/\sqrt{6} & 0 & 0 \end{bmatrix}, \quad (3.13)$$

where $\sin \theta_{12}^q \equiv \lambda = a_d - a_u$. We summarize some assumptions we used ; 1) Mass matrix of up quark and one of down quark are symmetric matrix. 2) Mass matrix M_u and M_d are the function of ϵ_u and ϵ_d , respectively. 3) Eq. (3.7) should hold for arbitrary ϵ_q .

It is straightforward to determine higher order corrections in our approach.

4 Comment on Lepton sector

Here we comment on lepton sector. In the democratic symmetry limit, the predicted lepton flavor mixing angles are given in eq. (2.12). If the symmetry breaking terms of charged leptons are the same as the ones of down quarks¹, we can estimate the correction

¹ We expect the symmetry breaking terms of down quark mass matrix can be applied for the ones of charged leptons. This is one of the reasons why we concentrate on the quark sector in this letter.

from zero-th order prediction, i.e., eq. (2.12). Note here that neutrinos are symmetry limit. In this case, lepton mixing angles are predicted as follows²,

$$\sin^2 2\theta_{sol} \sim 0.97, \quad \sin^2 2\theta_{atm} \sim 0.88, \quad U_{e3} \sim 0.18. \quad (4.1)$$

These predictions are slightly different from experimental data. This facts might indicate that we should consider $O(\lambda^2)$ corrections in CKM mixing matrix and/or the off-diagonal elements *and* small symmetry breaking terms of neutrino sector.

Then we present the estimation of the lepton flavor mixing angles when we include $O(\lambda^2)$ corrections in CKM mixing matrix (Note here that neutrinos are also symmetry limit.),

$$\sin^2 2\theta_{sol} \sim 0.87, \quad \sin^2 2\theta_{atm} \sim 0.80, \quad U_{e3} \sim 0.18. \quad (4.2)$$

Note that there is no effects for U_{e3} from $O(\lambda^2)$ corrections. Although $O(\lambda^3)$ terms in the unitary matrix V_l can modify U_{e3} , this effects would be neumerically too small to be consistent with experimental upper bound $U_{e3} < 0.15$ [4]. Thus the off-diagonal elements and the symmetry breaking terms in neutrino sector might be very important to obtain the realistic lepton mixing angles. In other words, small r parameter³ in eq. (2.10) *and* small symmetry breaking terms [9] are required.

5 Conclusion

In this letter, we investigate the symmetry breaking pattern of the democratic mass matrix model. We present the symmetry breaking terms in quark sector which are determined by alternative ways instead of conventional ansatz. These symmetry breaking terms might be useful for understanding the origin of democratic symmetry and the mechanism of its breaking. We also estimate the lepton fravor mixing angles by assuming that the symmetry breaking terms of charged leptons are the same as the ones of down quarks. The results indicate that to aquire realistic lepton mixing angles, the small

²Here we assumed $a_d \gg a_u$ and this leads to $a_d \sim O(\lambda)$. This reason is as follows. Since the unitary matrix V_q should become $V_q^{[0]}$ in the $\epsilon_q^2 \rightarrow 0$ limit, a_q (parameter in the unitary matrix V_q) might be proportional to $(\epsilon_q^2)^x$ (where x is positive unknown parameter). Thus λ might be expressed by $\lambda = (\epsilon_d^2)^x a'_d - (\epsilon_u^2)^x a'_u$. This relation is suitful for well-known phenomenological relation [11], $\lambda \sim \sqrt{m_d/m_s}$.

³ It is interesting that higher dimensional GUTs model which leads this situation is recently proposed [12].

off-diagonal elements *and* the symmetry breaking terms in neutrino mass matrix are important.

Acknowledgments

We would like to thank T. Inagaki, T. Onogi, K. Ohkura and M. Tanimoto for useful discussions.

References

- [1] H. Harari, H. Haut and J. Weyers, Phys. Lett. B **78** (1978) 459.
- [2] Y. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **81** (1998) 1158 [Erratum-ibid. **81** (1998) 4279] [hep-ex/9805021];
Y. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **81** (1998) 1562 [hep-ex/9807003];
Y. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **86** (2001) 5656 [hep-ex/0103033].
- [3] Q. R. Ahmad *et al.* [SNO Collaboration], hep-ex/0204008.
- [4] M. Apollonio *et al.* [CHOOZ Collaboration], Phys. Lett. B **466** (1999) 415 [hep-ex/9907037]
- [5] Y. Koide, Phys. Rev. D **28** (1983) 252; Phys. Rev. D **39** (1989) 1391;
M. Tanimoto, Phys. Rev. D **41** (1990) 1586;
K. Matsumoto, Prog. Theo. Phys. **84** (1990) 185;
H. Fritzsch and J. Plankl, Phys. Lett. B **237** (1990) 451.
- [6] Using alternative approach,
P. Kaus and S. Meshkov, Mod. Phys. Lett. A **3** (1988) 1251.
- [7] M. Fukugita, M. Tanimoto, T. Yanagida, Phys. Rev. D **57** (1998) 4429 [hep-ph/9709388].
- [8] H. Fritzsch and Z. Z. Xing, Phys. Lett. B **440** (1998) 313 [hep-ph/9808272].
- [9] M. Fujii, K. Hamaguchi, T. Yanagida, hep-ph/0202210.

- [10] H. Fritzsch and Z. Z. Xing, Prog. Part. Nucl. Phys. **45** (2000) 1 [hep-ph/9912358] and references there in.
- [11] For example, M. Tanimoto, hep-ph/9807517.
- [12] T. Watari and T. Yanagida, hep-ph/0205090.